arXiv:1210.5550v2 [astro-ph.CO] 23 Oct 2012

Tachyonic (phantom) power-law cosmology

Rachan Rangdee[1] and Burin Gumjudpai*[1, 2, †]
[1] *ThEP's CRL, NEP, The Institute for Fundamental Study,*
Naresuan University, Phitsanulok 65000, Thailand
[2] *Thailand Center of Excellence in Physics, Ministry of Education, Bangkok 10400, Thailand*
(Dated: October 24, 2012)

Tachyonic scalar field-driven late universe with dust matter content is considered. The cosmic expansion is modeled with power-law and phantom power-law expansion at late time ($z < 2$ for power-law and $z < 0.45$ for phantom power-law). WMAP7 and its combined data are used to constraint the model. For quintessence and tachyonic field, forms of the equation of state do not depends on type of the scalar field but depend only on scale factor function. However the forms of potential and the field solution are different in quintessence and tachyonic cases. Power-law cosmology model (driven with either quintessence or tachyonic field) predicts equation of state unmatched to the observational value, hence the model is excluded for quintessence and tachyonic field. The phantom power-law cosmology model predicts values of equation of state which agree with observational data (true for both quintessence and tachyonic cases), i.e. $w_{\phi,0} = -1.49^{+11.64}_{-4.08}$ (WMAP7+BAO+H_0) and $w_{\phi,0} = -1.51^{+3.89}_{-6.72}$ (WMAP7). The phantom-power law exponent β must be less than about -6, so that the $-2 < w_{\phi,0} < -1$. The phantom power-law tachyonic potential is reconstructed. The dimensionless potential slope variable Γ at present is about 1.5. The potential reduced to $V = V_0\phi^{-2}$ in the limit $\Omega_{\mathrm{m},0} \to 0$.

PACS numbers: 98.80.Cq

I. INTRODUCTION

There have been clear evidences that the present universe is under accelerating expansion as being observed by various observations, e.g. the cosmic microwave background (CMB) [1–3], large-scale structure surveys [4], supernovae type Ia (SNIa) [5–9] and X-ray luminosity from galaxy clusters [10]. One prime explanation is that the acceleration is an effect of a scalar field which could evolve under its potential to have negative pressure with $p < -\rho c^2/3$ giving repulsive gravity. Form of energy with this negative pressure range is generally called dark energy [11]. It is known that scalar field is responsible for symmetry breaking mechanisms and super-fast expansion in inflationary scenario, resolving horizon and flatness problems as well as explaining the origin of structures [12]. Introducing a cosmological constant into the field equation is simplest way to have dark energy [13], however it creates new problem on fine-tuning of energy density scales [14]. For the cosmological constant to be viable, idea of varying cosmological constant needs to be considered [15]. Considering dark energy as scalar field, there is possibility that the field could have non-canonical kinetic part such as tachyon which is classified as a type of k-essence models [16]. The tachyon field is a negative mass mode of unstable non-BPS D3-brane in string theory [17] or a massive scalar field on anti-D3 brane [18]. It was found that the tachyonic field potential must not be too steep, i.e. less steep than $V(\phi) \propto \phi^{-2}$ in order to account for the late acceleration [19] [20].

In this work, we considered dark energy in form of tachyonic scalar field in power-law cosmology whose scale factor scaled as $a \propto t^{\alpha}$ with $0 \leq \alpha \leq \infty$. The power law can give acceleration when $\alpha > 1$. In history of the universe, there were epoch when radiation or dust is dominant component in the universe for which the scale factor evolves as power-law $a \propto t^{1/2}$ and $a \propto t^{2/3}$. A universe with mixed combination of different cosmic ingredients can be modeled using power-law expansion with some approximately constant α during not-so-large period of cosmic time. Adjustability of expansion rate is characterized by only one parameter, α which is used widely in astrophysical observations. Non-minimally coupled scalar-tensor theory in which the scalar field couples to the curvature contributing to energy density that cancels out the vacuum energy can motivate the power-law cosmology [21]. In simple inflationary model, the power-law cosmology can avoid flatness and horizon problems and can give simple spectrum [22]. It has proved to be a very good phenomenological description of the cosmic evolution, since it can describe radiation epoch, dark matter epoch, and dark energy epoch according to value of the exponent [23, 24]. Previously linear-coasting cosmology,

---

* Corresponding author
[†] Electronic address: `buring@nu.ac.th`

$\alpha \approx 1$ was analyzed [25–28] with motivation from SU(2) instanton cosmology [29], higher order (Weyl) gravity [30], or from scalar-tensor theories [33]. However the universe expanding with $\alpha = 1$ [31] was not able to agree with observational constraint from Type Ia supernovae, Hubble rate data from cosmic chronometers and BAO [32] which indicates requirement that $H'(z)$=const and $q(z) = 0$ are not favored by the observations.

For a specific gravity or dark energy model, power-law cosmology is considered in $f(T)$ and $f(G)$ gravities [34] and in the case of which there is coupling between cosmic fluids [35]. The power-law cosmology were also studied in context of scalar field cosmology [36, 37], phantom scalar field cosmology [38]. There is also slightly different form of the power-law function which α can evolved with time so that it can parameterize cosmological observables [39].

For the power-law to be valid throughout the cosmic evolution, it is not possible with constant exponent. For example, at big bang primordial nucleosynthesis (BBN), α is allowed to have maximum value at approximately 0.55 in order to be capable of light element abundances [40, 41]. The value is about 1/2 at highly-radiation dominated era, about 2/3 at highly-dust dominated era and greater than one at present. Low value of α results in much younger cosmic age and does not give acceleration. On the other hand $\alpha \geq 1$ value is needed to solve age problem in the CDM model [23] without flatness and horizon problems. In universe dominated with cold dark matter and dark energy, considering that the power-law expansion happens long after matter-radiation equality era, $z \ll 3196$ (value from [2]), the BBN constraint can be relaxed and large α is allowed. We consider power-law cosmology with a small range of recent cosmic era when dark energy began to dominate, i.e. from $z \lesssim 2$ to present using results from WMAP7 [2] and WMAP7+BAO+H_0 combined datasets [3]. There are tachyonic scalar field evolving under potential $V(\phi)$ and dust barotropic fluid (cold dark matter and baryonic matter) as two major ingredients. We aim to test whether the power-law cosmology is still valid in the scenario of tachyonic scalar field by looking at value of the equation of state predicted by the power-law tachyonic cosmology and that of model-independent direct-observational result. The WMAP7 and WMAP7+BAO+H_0 data used here are presented in Table I. We also consider when the field is phantom, i.e. having negative kinetic term with phantom-power law expansion [42], $a \propto (t_{\rm s} - t)^{\beta}$, $\beta < 0$ from $z \lesssim 0.45$ till present. We determine tachyonic field equation of state parameter, w_ϕ and we perform parametric plot versus exponent β. We then analyze the result and conclude this work.

II. BACKGROUND COSMOLOGY AND OBSERVATIONAL DATA

We consider standard FLRW universe containing dust matter (cold dark matter and baryonic matter) with tachyonic field with Lagrangian,

$$\mathcal{L}_{\rm tachyon} \;=\; -V(\phi)\sqrt{1-\partial_\mu\phi\partial^\mu\phi} \tag{1}$$

evolving under the background Friedmann equation,

$$H^2 = \frac{8\pi G}{3}\left(\rho_\phi + \rho_{\rm m}\right) - \frac{kc^2}{a^2} \tag{2}$$

and acceleration rate,

$$\dot{H} = \frac{\ddot{a}}{a} - H^2 = -\frac{4\pi G}{c^2}\left(\rho_\phi c^2 + p_\phi + \rho_{\rm m}c^2 + p_{\rm m}\right) + \frac{kc^2}{a^2} \tag{3}$$

Tachyonic field energy density and pressure are

$$\rho_\phi c^2 \;=\; \frac{V(\phi)}{\sqrt{1-\epsilon\dot{\phi}^2}} \tag{4}$$

$$p_\phi \;=\; -V(\phi)\sqrt{1-\epsilon\dot{\phi}^2} \tag{5}$$

where $\epsilon = \pm 1$. The negative ϵ represents the case when kinetic term of the tachyon is phantom. The tachyonic fluid equation reads

$$\frac{\epsilon\ddot{\phi}}{1-\epsilon\dot{\phi}^2} + 3H\epsilon\dot{\phi} + \frac{V'}{V} = 0 \tag{6}$$

Using Eq. (4), (5) and (6) in the (3) (dust pressure is zero), we obtain

$$\dot{H} = -\frac{4\pi G}{c^2}\left(\frac{V\epsilon\dot{\phi}^2}{\sqrt{1-\epsilon\dot{\phi}^2}} + \rho_{\rm m}c^2\right) + \frac{kc^2}{a^2} \tag{7}$$

Using tachyonic density (4) in the Friedmann equation therefore

$$\frac{V}{\sqrt{1-\epsilon\dot{\phi}^2}} = \frac{3}{8\pi G/c^2}\left(H^2+\frac{kc^2}{a^2}\right) - \rho_{\rm m}c^2 \tag{8}$$

Substituting (8) into the equation (7), we obtain

$$\dot{H} = -4\pi G\left[\frac{3\epsilon\dot{\phi}^2}{8\pi G}\left(H^2+\frac{kc^2}{a^2}\right) - \rho_{\rm m}\epsilon\dot{\phi}^2 + \rho_{\rm m}\right] + \frac{kc^2}{a^2} \tag{9}$$

which can be rewritten as

$$\epsilon\dot{\phi}^2 \;=\; -\left[\frac{2\dot{H}-(2kc^2/a^2)+8\pi G\rho_{\rm m}}{3H^2+(3kc^2/a^2)-8\pi G\rho_{\rm m}}\right] \tag{10}$$

and hence

$$1-\epsilon\dot{\phi}^2 \;=\; \frac{3H^2+2\dot{H}+(kc^2/a^2)}{3H^2+(3kc^2/a^2)-8\pi G\rho_{\rm m}} \tag{11}$$

We use the above expression in the equation (8), as a result we can get tachyonic potential

$$V \;=\; \left[\frac{3}{8\pi G/c^2}\left(H^2+\frac{kc^2}{a^2}\right) - \rho_{\rm m}c^2\right]\sqrt{\frac{3H^2+2\dot{H}+(kc^2/a^2)}{3H^2+(3kc^2/a^2)-8\pi G\rho_{\rm m}}}. \tag{12}$$

The tachyonic equation of state, w_ϕ is, from (11),

$$w_\phi \;=\; \frac{p}{\rho c^2} \;=\; -(1-\epsilon\dot{\phi}^2) \;=\; -\left[\frac{3H^2+2\dot{H}+(kc^2/a^2)}{3H^2+(3kc^2/a^2)-8\pi G\rho_{\rm m}}\right] \tag{13}$$

This can be weighed with the dust-matter content to give an effective equation of state, $w_{\rm eff} = \rho_\phi w_\phi/(\rho_\phi+\rho_{\rm m})$. With all information above, $w_{\rm eff}$ is expressed as

$$w_{\rm eff} \;=\; w_\phi\left[1-\frac{8\pi G\rho_{\rm m}/3}{H^2+(kc^2/a^2)}\right] \tag{14}$$

We found that the equation (13) and (14) are the same for both quintessence scalar field [37] and tachyonic field cases, albeit the $\dot{\phi}$ and $V(\phi)$ are expressed differently in both cases. That is for both quintessence and tachyonic cases, w_ϕ does not depend on the scalar field model but depends on the form of expansion function. This is also true for $w_{\rm eff,0}$. The equation of state is also independent of the sign of ϵ which indicates negative kinetic energy. Using power-law expansion and phantom power-law expansion into (13), one can find the present value of the equation of state, $w_{\phi,0}$. This value is a (phantom) power-law prediction of the $w_{\phi,0}$. We can compare this predicted value to the model-independent $w_{\phi,0}$ obtained from CMB observation.

The derived data from WMAP7+BAO+H_0 and WMAP7 are presented in Table I. We will set $a_0 = 1$ and consider flat universe $k = 0$ (but keep it in formulae for completeness). Dust density is defined as $\rho_{\rm m,0} = \Omega_{\rm m,0}\rho_{\rm c,0}$. Total dust fluid density at present is sum of that of all dust mater types $\Omega_{\rm m,0} = \Omega_{\rm CDM,0} + \Omega_{\rm b,0}$. Present value of the critical density is $\rho_{\rm c,0} = 3H_0^2/8\pi G$, and radiation density is negligible. We take the maximum likelihood value assuming spatially flat case. Although in deriving t_0, the ΛCDM model is assumed with the CMB data, however one can estimably use t_0 since $w_{\rm DE}$ is very close to -1. In SI units, the reduced Planck mass squared is $M_{\rm P}^2 = \hbar c/8\pi G$. In this work, we also give correction to errors on future singularity time, $t_{\rm s}$ (phantom power-law case) reported previously in [38] and improve values of $w_{\phi,0}$ of the phantom power-law case in [38] and of the usual power-law case reported earlier [37].

III. POWER-LAW COSMOLOGY

In power-law cosmology, scale factor is a function of time as

$$a(t) = a_0\left(\frac{t}{t_0}\right)^{\alpha}, \tag{15}$$

Parameter	**WMAP7+BAO+H_0**	**WMAP7**
t_0	13.76 ± 0.11 Gyr or $(4.34 \pm 0.03) \times 10^{17}$ sec	13.79 ± 0.13 Gyr or $(4.35 \pm 0.04) \times 10^{17}$ sec
H_0	70.4 ± 1.4 km/s/Mpc	70.3 ± 2.5 km/s/Mpc
	$(2.28 \pm 0.04) \times 10^{-18}$ sec^{-1}	$(2.28 \pm 0.08) \times 10^{-18}$ sec^{-1}
$\Omega_{\rm b,0}$	0.0455 ± 0.0016	0.0451 ± 0.0028
$\Omega_{\rm CDM,0}$	0.226 ± 0.015	0.226 ± 0.027
$\Omega_{\rm m,0}$	$0.271(5) \pm 0.015(1)$	$0.271(1) \pm 0.027(1)$
$\rho_{\rm m,0}$	$(2.52(49)^{+0.18(24)}_{-0.16(64)}) \times 10^{-27}$ kg/m^3	$(2.52(12)^{+0.30(97)}_{-0.30(61)}) \times 10^{-27}$ kg/m^3
$\rho_{\rm c,0}$	$(9.29(99)^{+0.32(92)}_{-0.32(35)}) \times 10^{-27}$ kg/m^3	$(9.29(99)^{+0.66(41)}_{-0.64(12)}) \times 10^{-27}$ kg/m^3

TABLE I: Combined WMAP7+BAO+H_0 and WMAP7 derived parameters (maximum likelihood) from Refs. [2] and [3]. Here we also calculate (with error analysis) $\Omega_{\rm m,0} = \Omega_{\rm b,0} + \Omega_{\rm CDM,0}$, critical density: $\rho_{\rm c,0} = 3H_0^2/8\pi G$ and matter density: $\rho_{\rm m,0} = \Omega_{\rm m,0}\rho_{\rm c,0}$. The space is flat and a_0 is set to unity.

The speed is $\dot{a} = \alpha a/t$ and the acceleration is $\ddot{a} = \alpha(\alpha - 1)a/t^2$. The Hubble parameter is $H(t) = \dot{a}/a = \alpha/t$ with $\dot{H} = -\alpha/t^2$. The deceleration parameter in this scenario is $q \equiv -a\ddot{a}/\dot{a}^2 = (1/\alpha) - 1$, that is $\alpha = 1/(q+1)$. As $\alpha \geq 0$ is required in power-law cosmology, hence $q \geq -1$ and $H_0 \geq 0$. To convert into redshift z, from $1 + z = a_0/a$ then $1 + z = (t_0/t)^\alpha$. Typically astrophysical tests for power-law cosmology indicating the value of α are performed by observing $H(z)$ data of SNIa or high-redshift objects such as distant globular clusters [43–45]. To indicate the value of α one can also use gravitational lensing statistics [28], compact-radio source [46] or using X-ray gas mass fraction measurements of galaxy clusters [47, 48]. Study of angular size to z relation of a large sample of milliarcsecond compact radio sources in flat FLRW universe found that $\alpha = 1.0 \pm 0.3$ at 68 % C.L. [46]. WMAP5 dataset gives $\alpha = 1.01$ for closed geometry [36]. Some procedures of measurement give large value of α such as $\alpha = 2.3^{+1.4}_{-0.7}$ (X-ray mass fraction data of galaxy clusters in flat geometry) [47] and $\alpha = 1.62^{+0.10}_{-0.09}$ (joint test using Supernova Legacy Survey (SNLS) and $H(z)$ data in flat geometry) [43]. Notice that assumption of spatial curvature is assumed in these results in the process of finding α except the WMAP5 which possesses constraint on spatial curvature in its results. When α is found with curvature-independent procedure (i.e. with neither SNIa nor cluster X-ray gass mass fraction) or in flat case, α value is near unity. For example, $H(z)$ data gives $\alpha = 1.07^{+0.11}_{-0.09}$ [43] and $\alpha = 1.11^{+0.21}_{-0.14}$ [37, 45]. Short review of recent α values can be found in Ref. [37]. Here α is calculated from value at present H_0, t_0 as $\alpha = H_0 t_0$. From (13) and (14), in case of power-law cosmology driven by tachyonic field, the equation of state of dark energy is

$$w_\phi = -\left[\frac{(3\alpha^2/t^2) - (2\alpha/t^2) + (kc^2/a_0^2)(t_0/t)^{2\alpha}}{(3\alpha^2/t^2) + (3kc^2/a_0^2)(t_0/t)^{2\alpha} - 8\pi G\rho_{\rm m,0}(t_0/t)^{3\alpha}}\right] \tag{16}$$

and

$$w_{\rm eff} = w_\phi\left[1 - \frac{(8\pi G/3)\rho_{\rm m,0}(t_0/t)^{3\alpha}}{(\alpha^2/t^2) + (kc^2/a_0^2)(t_0/t)^{2\alpha}}\right] \tag{17}$$

At present, $t = t_0$, $w_{\rm eff,0} = -1 + 2/(3\alpha)$. In Table II, values of equation of state parameters derived in the power-law cosmology (true for both tachoynic and quintessence) do not match observational data, i.e. $w_{\phi,0}$ and $w_{\rm eff,0}$ found here are much greater than observational (spatially flat) WMAP derived results, for example WMAP7 data[1]: $w_{\phi,0} = -1.12^{+0.42}_{-0.43}$, WMAP7+BAO+$H_0$ combined data[2]: $w_{\phi,0} = -1.10^{+0.14}_{-0.14}$ (68 % CL), WMAP7+BAO+H_0+SN data[3]: $w_{\phi,0} = -1.34^{+1.74}_{-0.36}$ (68 % CL) and WMAP7+BAO+H_0+SN data with time delay distance information correction[4]: $w_{\phi,0} = -1.31^{+1.67}_{-0.38}$ (68 % CL). We conclude that the quintessential scalar field [37] and tachyonic field in power-law expansion universe are not viable.

---

[1] flat geometry, constant $w_{\phi,0}$ (Sec. 4.2.5 of Ref. [2])

[2] flat geometry, constant $w_{\phi,0}$ (Sec. 5.1 of Ref. [3])

[3] flat geometry, time varying dark energy EoS, $w_\phi(a) = w_0 + w_a(1-a)$ with $w_0 = -0.93 \pm 0.13, w_a = -0.41^{+0.72}_{-0.71}$ (Sec. 5.3 of Ref. [3])

[4] flat geometry, time varying dark energy EoS, $w_\phi(a) = w_0 + w_a(1-a)$ with $w_0 = -0.93 \pm 0.12, w_a = -0.38^{+0.66}_{-0.65}$ (Sec. 5.3 of Ref. [3])

Parameter	WMAP7+BAO+H_0	WMAP7
α	$0.98(95) \pm 0.01(87)$	$0.99(18) \pm 0.03(60)$
$w_{\phi,0}$ (with power-law cosmology)	$-0.44(79)^{+0.01(66)}_{-0.01(54)}$	$-0.44(98)^{+0.02(97)}_{-0.02(82)}$
$w_{\rm eff,0}$ (with power-law cosmology)	$-0.32(63) \pm 0.01(25)$	$-0.32(78) \pm 0.02(35)$

TABLE II: Power-law cosmology exponent and its prediction of equation of state parameters. The value does not match the WMAP7 model-independent results.

IV. PHANTOM POWER-LAW COSMOLOGY

In this section, we can check if phantom power-law could be a valid solution for the tachyonic-driven universe (also true for quintessence). The phantom power law reads

$$a(t) = a_0 \left(\frac{t_{\rm s} - t}{t_{\rm s} - t_0} \right)^{\beta} \tag{18}$$

with speed, $\dot{a} = -a_0 \beta (t_{\rm s} - t)^{\beta - 1}/(t_{\rm s} - t_0)^{\beta} = -\beta a/(t_{\rm s} - t)$ and acceleration, $\ddot{a} = a_0 \beta(\beta - 1)\,(t_{\rm s} - t)^{\beta - 2}/(t_{\rm s} - t_0)^{\beta} = \beta(\beta - 1) a/(t_{\rm s} - t)^2$ where $t_{\rm s}$ is future big-rip singularity time [42]. For both $\dot{a}$ and $\ddot{a}$ to be greater than zero, i.e. both expanding and accelerating, the condition $\beta < 0$ is needed. The Hubble parameter is therefore,

$$H \;=\; \frac{\dot{a}}{a} \;=\; -\frac{\beta}{t_{\rm s} - t} \qquad \text{hence} \qquad \dot{H} \;=\; -\frac{\beta}{(t_{\rm s} - t)^2} \tag{19}$$

At present, we can find $\beta = H_0 (t_0 - t_{\rm s})$. The deceleration parameter is $q \;\equiv\; -a\ddot{a}/\dot{a}^2 \;=\; (1/\beta) - 1$. The dust matter density, $\rho_{\rm m} = \rho_{\rm m,0} a_0^3/a^3$ is then

$$\rho_{\rm m} = \rho_{\rm m,0} \left(\frac{t_{\rm s} - t_0}{t_{\rm s} - t} \right)^{3\beta} \tag{20}$$

Substituting these equations into (13) and (14), we obtain,

$$w_{\phi} \;=\; -\left[\frac{\beta(3\beta - 2)/(t_{\rm s} - t)^2 \;+\; (kc^2/a_0^2)\left[(t_{\rm s} - t_0)/(t_{\rm s} - t)\right]^{2\beta}}{3\beta^2/(t_{\rm s} - t)^2 \;+\; (3kc^2/a_0^2)\left[(t_{\rm s} - t_0)/(t_{\rm s} - t)\right]^{2\beta} \;-\; 8\pi G \rho_{\rm m,0}\left[(t_{\rm s} - t_0)/(t_{\rm s} - t)\right]^{3\beta}} \right] \tag{21}$$

$$w_{\rm eff} \;=\; w_{\phi} \left[1 - \frac{(8\pi G/3)\rho_{\rm m,0}\left[(t_{\rm s} - t_0)/(t_{\rm s} - t)\right]^{3\beta}}{\beta^2/(t_{\rm s} - t)^2 \;+\; (kc^2/a_0^2)\left[(t_{\rm s} - t_0)/(t_{\rm s} - t)\right]^{2\beta}} \right] \tag{22}$$

To convert to redshift one can use $1 + z = a_0/a$ therefore $1 + z = \left[(t_{\rm s} - t_0)/(t_{\rm s} - t)\right]^{\beta}$ and $t_{\rm s} - t = (t_{\rm s} - t_0)(1 + z)^{-1/\beta}$. At present, $t = t_0$, $w_{\rm eff,0} = -1 + 2/(3\beta)$. The big-rip time $t_{\rm s}$, can be calculated from

$$t_{\rm s} \approx t_0 - \frac{2}{3(1 + w_{\rm DE})} \frac{1}{H_0 \sqrt{1 - \Omega_{\rm m,0}}} \tag{23}$$

Here, $w_{\rm DE}$ must be less than -1 and in deriving this above expression flat geometry and constant dark energy equation of state is assumed [42]. We will estimably use $t_{\rm s}$ from this formula. In finding error bar of $t_{\rm s}$, we exploit better procedure than that performed earlier in [38] by considering that the second order of error bar multiplications are too large to approximate to zero. We discuss this in the appendix. Results presented in Table III are $\beta, t_{\rm s}$ and the equation of state. For phantom power-law cosmology driven by tachyonic field (also true for phantom quintessence), the resulting value is $w_{\phi,0} = -1.49^{+11.64}_{-4.08}$ (using WMAP7+BAO+H_0) and $-1.51^{+3.89}_{-6.72}$ (using WMAP7). These do not much differ from results from WMAP7+BAO+H_0+SN data (flat, varying dark energy EoS) which gives $w_{\phi,0} = -1.34^{+1.74}_{-0.36}$ (68 % CL) and WMAP7+BAO+H_0+SN+time delay distance correction data (flat varying dark energy EoS) which gives $w_{\phi,0} = -1.31^{+1.67}_{-0.38}$ (68 % CL) [3]. Using observational data in Tables I and III we derive

$$w_{\phi,0} \;=\; -\left[\frac{1 - 2/(3\beta)}{1 - (16.60/\beta^2)} \right] \qquad (\text{WMAP7} + \text{BAO}{+}H_0) \tag{24}$$

$$w_{\phi,0} \;=\; -\left[\frac{1 - 2/(3\beta)}{1 - (11.47/\beta^2)} \right] \qquad (\text{WMAP7}) \tag{25}$$

With these, we show parametric plots of the $w_{\phi,0}$ and β in Fig. 1. The values measured for β and $w_{\phi,0}$ are the purple cross (WMAP7+BAO+H_0) and yellow spot (WMAP7). For $-\infty < \beta \lesssim -6$, $w_{\phi,0}$ lies in the range $(-1,-2)$. Fig. 2 shows evolution of $w(z)$ in late phantom power-law universe from $0 < z < 0.45$, i.e. $t = 8.48$ Gyr (both datasets) till present era (this is to avoid singularity in w_ϕ at $z = 0.492$ (WMAP7+BAO+H_0) and at $z = 0.484$ (WMAP7)). These are equivalent to 5.28 Gyr ago (WMAP7+BAO+H_0) and 5.31 Gyr ago (WMAP7).

Parameter	**WMAP7+BAO+H_0**	**WMAP7**
β	$-7.81(08)^{+11.71(8)}_{-4.56(1)}$	$-6.50(72)^{+3.91(92)}_{-5.09(96)}$
t_s	$122.30(0)^{+162.83(7)}_{-63.36(0)}$ Gyr	$104.21(5)^{+54.37(3)}_{-70.79(9)}$ Gyr
$w_{\phi,0}$ (with phantom power-law cosmology)	$-1.48(99)^{+11.64(46)}_{-4.08(45)}$	$-1.51(26)^{+3.89(23)}_{-6.71(90)}$
$w_{\mathrm{eff},0}$ (with phantom power-law cosmology)	$-1.08(54)^{+0.25(60)}_{-0.11(98)}$	$-1.10(24)^{+0.15(52)}_{-0.37(12)}$

TABLE III: Phantom power-law cosmology exponent and its prediction of equation of state parameters. The equation of state lies in acceptable range of values given by WMAP7 results. Large error bar of $w_{\phi,0}$ is an effect of large error bar in t_s.

V. TACHYONIC POTENTIAL FOR PHANTOM POWER-LAW COSMOLOGY

From (12) the tachyonic potential of the phantom-power law is not the same as the quintessential potential. We start from considering equation (10) for flat space and $\epsilon = -1$ hence $\dot{\phi}^2 \; = \; (2\dot{H} + 8\pi G\rho_\mathrm{m})/(3H^2 - 8\pi G\rho_\mathrm{m})$. We make approximation that the dust term is much less contributive compared to the $\dot{H}$ and H^2 terms therefore,

$$\dot{\phi}^2 \;\approx\; \frac{2\dot{H}}{3H^2} \;=\; -\frac{2}{3\beta} \tag{26}$$

Since $\beta < 0$ always then $-\beta = |\beta|$. Integrating from t to t_s, and choosing positive solution,

$$\phi(t) \;\approx\; \sqrt{\frac{2}{3|\beta|}}\;(t_\mathrm{s} - t) \tag{27}$$

We set $B \equiv \sqrt{3|\beta|/2}$, hence $t_\mathrm{s} - t = B\phi$. Using the equation (12) we find that

$$V(\phi) \approx \left[\frac{3c^2\beta^2}{\kappa(B\phi)^2} - \rho_{\mathrm{m},0}\, c^2 \left(\frac{t_\mathrm{s} - t_0}{B\phi}\right)^{3\beta}\right] \times \left[\frac{1 - 2/(3\beta)}{1 - \rho_{\mathrm{m},0}[\kappa/(3\beta^2)]\,(B\phi)^2}\right]^{1/2} \tag{28}$$

where $\kappa \equiv 8\pi G$. Note that the term $1 - 2/(3\beta)$ is just $-w_{\mathrm{eff},0}$. We can rearrange the potential in form of cosmological observables $H_0, \Omega_{\mathrm{m},0}$ and q,

$$V(\phi) \approx \frac{c^2}{\kappa}\left[\frac{2|\beta|}{\phi^2} \;-\; 3\left(\frac{3}{2}\right)^{3|\beta|/2} |\beta|^{-3|\beta|/2}\,\Omega_{\mathrm{m},0}\; H_0^{2+3|\beta|}\;\phi^{3|\beta|}\right] \times \left[\frac{1 + 2/(3|\beta|)}{1 - (3/2)\,\Omega_{\mathrm{m},0}\,H_0^2\phi^2/|\beta|}\right]^{1/2} \tag{29}$$

where $\beta = \beta(q) = (1+q)^{-1}$. This is plotted in Fig. 3 where the field values at present and at $z = 0.45$ are

$$\begin{array}{llllll} \phi(z=0) = 1.268 \times 10^{17} & \text{sec} & \text{and} & \phi(z=0.45) = 7.803 \times 10^{16} & \text{sec} & (\text{WAMP7} + \text{BAO} + H_0) \\ \phi(z=0) = 1.392 \times 10^{17} & \text{sec} & \text{and} & \phi(z=0.45) = 8.555 \times 10^{16} & \text{sec} & (\text{WAMP7}) \end{array}$$

It has been known that in order for the tachyonic potential to account for the late acceleration, it should not be steeper than the potential $V \propto \phi^{-2}$ [19]. To check if our derived tachyonic potential could fit in this criteria, that is shallower than $V \propto \phi^{-2}$, we use dimensionless variable,

$$\Gamma = \frac{V''V}{V'^2} \tag{30}$$

where $'$ denotes $\mathrm{d}/\mathrm{d}\phi$. For the potential $V \propto \phi^{-2}$, we have $\Gamma = 3/2$. Hence in general the potential with $\Gamma < 3/2$ satisfies this criteria. Considering the potential (29) we use both derived datasets to compute its dynamical slope $\Gamma(\phi)$



FIG. 1: Present value of phantom tachyonic dark energy equation of state plotted versus β and their error bar resulting from error bar in β. This is the same in quintessence case.

which is in very complicated form. We plot this in Fig. 4. We found that using our data with the field value at present, for WMAP7+BAO+H_0, $\Gamma(\phi(z=0)) = 1.503$ and for WMAP7, $\Gamma(\phi(z=0)) = 1.505$. These values are approximately the same as that of $V \propto \phi^{-2}$. Note that the $V \propto \phi^{-2}$ potential is found when the universe is filled with tachyon field as single component. Indeed in the limit $\Omega_{\mathrm{m},0} \rightarrow 0$, our derived potential (29) becomes $V \propto \phi^{-2}$. Note that for other tachyonic potentials such as $V = V_0/[\cosh(a\phi/2)]$ and $V = V_0 e^{(1/2)m^2\phi^2}$, their slopes are $\Gamma = 1 - \mathrm{csch}^2(a\phi/2)$ and $1 + (m\phi)^{-2}$ respectively. These examples are typical tachyonic potentials which also have dynamical slopes. In Fig. 4, $\Gamma(\phi)$ diverges twice however, in the region we consider ($z = 0.45 \rightarrow z = 0$), the value of Γ stays approximately at 1.5.

VI. CONCLUSION

In this work model of tachyonic-driven universe are investigated for usual power-law cosmology and phantom power-law cosmology. The universe is flat FLRW filled with tachyonic scalar field and dust matter. We consider late universe when dark energy begins to dominate, i.e. $z < 2$ for power-law cosmology and for $z < 0.45$ for phantom power-law cosmology. WMAP7 data and its derived data when combined with BAO and $H(z)$ data are used in this study. We find power-law, phantom-power-law exponents and other cosmological observables and we improve data reported earlier in [38]. We find that although the forms of potential and the field solution are different for quintessential scalar field [37, 38] and tachyonic field, however the equation of state are in identical form for both quintessential scalar field and tachyonic field. This is to say that, for quintessence and tachyonic field, the equation of state does not depends on type of the scalar field but depends only on forms of expansion function of the scale factor. The present value of dark energy equation of state predicted by quintessential and tachyonic power-law cosmology models do not match both WMAP7 datasets. We conclude that the usual power-law cosmology model with either quintessence or with tachyonic field are excluded by these observational data. When considering the other case, i.e the phantom power-law cosmology, the model predicts values of equation of state not much differ from results from observational data (true for both quintessence and tachyonic cases), i.e. $w_{\phi,0} = -1.49^{+11.64}_{-4.08}$ (phantom power-law using WMAP7+BAO+H_0) and $w_{\phi,0} = -1.51^{+3.89}_{-6.72}$ (phantom power-law using WMAP7 data) compared to $w_{\phi,0} = -1.34^{+1.74}_{-0.36}$ (WMAP7+BAO+H_0+SN data): and $w_{\phi,0} = -1.31^{+1.67}_{-0.38}$ (WMAP7+BAO+H_0+SN+time delay distance correction data) [3]. From parametric plot in Fig. 1, at $\beta \lesssim -6$, $w_{\phi,0}$ is in the expected range (-2, -1). We reconstruct the tachyonic potential in this scenario and we find that the dimensionless slope variable Γ of our derived potential at present time is about 1.5. The phantom-power-law tachyonic potential found here reduced to $V = V_0\phi^{-2}$



FIG. 2: Phantom tachyonic (and quintessence) dark energy equation of state versus z.



FIG. 3: Potential versus field using WMAP7+BAO+H_0, WMAP7

in the limit $\Omega_{\mathrm{m},0} \to 0$.

Acknowledgments

We thank Chakkrit Kaeonikhom for discussion. B. G. is sponsored by a project number: BRG5380018 under the Basic Research Grant of the Thailand Research Fund (TRF). R. R. is funded via the TRF's Royal Golden Jubilee



FIG. 4: Dimensionless variable Γ plotted versus field using WMAP7+BAO+H_0 and WMAP7. The considered region for late universe $z < 0.45$ lies in the bars.

Doctoral Scholarship.

Appendix: Errors Analysis

In calculating of the accumulated errors, we follow the procedure here. If f is valued of answer in the form

$$f = f(x_1, x_2, \dots, x_n) \tag{A.1}$$

and f_0 is the value when x_i is set to their measured values, then the value of f_i is defined as

$$f_i = f(x_1, \dots, x_i + \sigma_i, \dots, x_n) \tag{A.2}$$

This value of f is the value with effect of error in variable x_i, that is σ_i. One can find square of the accumulated error from

$$\sigma_f^2 = \sum_i^n (f_i - f_0)^2 \tag{A.3}$$

Hence giving the error of f from accumulating effect from errors of x_i.

[1] S. Masi *et al.*, Prog. Part. Nucl. Phys. **48**, 243 (2002).

[2] D. Larson, J. Dunkley, G. Hinshaw, E. Komatsu, M. R. Nolta, C. L. Bennett, B. Gold, M. Halpern *et al.*, Astrophys. J. Suppl. **192**, 16 (2011).

[3] E. Komatsu *et al.* (WMAP Collaboration), Astrophys. J. Suppl. **192**, 18 (2011).

[4] R. Scranton *et al.* (SDSS Collaboration), astro-ph/0307335; M. Tegmark *et al.* (SDSS Collaboration), Phys. Rev. D **69**, 103501 (2004).

[5] S. Perlmutter *et al.* (Supernova Cosmology Project Collaboration), Nature **391**, 51 (1998); S. Perlmutter *et al.* (Supernova Cosmology Project Collaboration), Astrophys. J. **517**, 565 (1999).

[6] A. G. Riess *et al.* (Supernova Search Team Collaboration), Astron. J. **116**, 1009 (1998); A. G. Riess, arXiv: astro-ph/9908237; G. Goldhaber *et al.* (The Supernova Cosmology Project Collaboration), Astrophys. J., **558**, 359 (2001); J. L. Tonry *et al.* (Supernova Search Team Collaboration), Astrophys. J. **594**, 1 (2003); A. G. Riess *et al.*, Astrophys. J. **659**, 98 (2007).

[7] A. G. Riess *et al.* (Supernova Search Team Collaboration), Astrophys. J. **607**, 665 (2004).

[8] P. Astier *et al.* (SNLS Collaboration), Astron. Astrophys. **447**, 31 (2006).

[9] R. Amanullah *et al.*, Astrophys. J. **716**, 712 (2010).

[10] S. W. Allen, R. W. Schmidt, H. Ebeling, A. C. Fabian and L. van Speybroeck, Mon. Not. Roy. Astron. Soc. **353**, 457 (2004); D. Rapetti, S. W. Allen and J. Weller, Mon. Not. Roy. Astron. Soc. **360**, 555 (2005).

[11] T. Padmanabhan, Curr. Sci. **88**, 1057 (2005); E. J. Copeland, M. Sami and S. Tsujikawa, Int. J. Mod. Phys. D **15**, 1753 (2006); T. Padmanabhan, AIP Conf. Proc. **861**, 179 (2006).

[12] A. A. Starobinsky, Phys. Lett. B **91**, 99 (1980); A. H. Guth, Phys. Rev. D **23**, 347 (1981); K. Sato, Mon. Not. Roy. Astro. Soc. **195**, 467 (1981); A. Albrecht and P. J. Steinhardt, Phys. Rev. Lett. **48**, 1220 (1982); A. D. Linde, Phys. Lett. B **108**, 389 (1982).

[13] S. Weinberg, Rev. Mod. Phys. **61**, 1 (1989); L. H. Ford, Phys Rev D **35**, 2339 (1987); A.D. Dolgov, Phys. Rev. D **55**, 5881 (1997).

[14] V. Sahni and A. Starobinsky, Int. J. Mod. Phy. D **9**, 373 (2000); P. J. Peebles and B. Ratra, Rev. Mod. Phys. **75**, 559 (2003).

[15] J. Sola and H. Stefancic, Phys. Lett. B **624**, 147 (2005); I. L. Shapiro and J. Sola, Phys. Lett. B **682**, 105 (2009).

[16] C. Armendariz-Picon, V. F. Mukhanov and P. J. Steinhardt, Phys. Rev. Lett. **85**, 4438 (2000); C. Armendariz-Picon, V. F. Mukhanov and P. J. Steinhardt, Phys. Rev. D **63**, 103510 (2001).

[17] M. R. Garousi, Nucl. Phys. B **584**, 284 (2000); A. Sen, JHEP **0207**, 065 (2002); A. Sen, JHEP **0204**, 048 (2002).

[18] M. R. Garousi, M. Sami and S. Tsujikawa, Phys. Rev. D **70**, 043536 (2004).

[19] T. Padmanabhan, Phys. Rev. D **66**, 021301 (2002); J. S. Bagla, H. K. Jassal and T. Padmanabhan, Phys. Rev. D **67**, 063504 (2003);

[20] D. Kutasov and V. Niarchos, Nucl. Phys. B **666**, 56 (2003); L. R. W. Abramo and F. Finelli, Phys. Lett. B **575**, 165 (2003); J. M. Aguirregabiria and R. Lazkoz, Phys. Rev. D **69**, 123502 (2004); E. J. Copeland, M. R. Garousi, M. Sami and S. Tsujikawa, Phys. Rev. D **71**, 043003 (2005).

[21] A. D. Dolgov in *The Very Early Universe*, ed. G. Gibbons, S. W. Hawking and S. T. Tiklos (Cambridge University Press, 1982); L. H. Ford, Phys. Rev. D **35**, 2339 (1987); Y. Fujii and T. Nishioka, Phys. Rev. D **42** 361 (1990); A. D. Dolgov, Phys. Rev. D **55**, 5881 (1997).

[22] F. Lucchin and S. Matarrese, Phys. Rev. D **32**, 1316 (1985).

[23] E. W. Kolb, Astrophys. J. **344**, 543 (1989).

[24] P. J. E. Peebles, *Principles of Physical Cosmology*, Princeton University Press (1993).

[25] D. Lohiya, S. Mahajan, A. Mukherjee and A. Batra, arXiv:astro-ph/9606082.

[26] M. Sethi, A. Batra and D. Lohiya, Phys. Rev. D **60**, 108301 (1999).

[27] A. Dev, M. Sethi and D. Lohiya, Phys. Lett. B **504**, 207 (2001).

[28] A. Dev, M. Safonova, D. Jain and D. Lohiya, Phys. Lett. B **548**, 12 (2002).

[29] R. E. Allen, arXiv: astro-ph/9902042

[30] P. Manheim and D. Kazanas, Gen. Rel. Grav. **22**, 289 (1990).

[31] F. Melia and A. Shevchuk, Mon. Not. Roy. Astron. Soc. **419**, 2579 (2012)

[32] M. Bilicki and M. Seikel, Mon. Not. Roy. Astron. Soc. **425**, 1664 (2012)

[33] D. Lohiya and M. Sethi, Class. Quant. Grav. **16**, 1545 (1999).

[34] A. R. Rastkar, M. R. Setare and F. Darabi, Astrophys. Space Sci. **337**, 487 (2012); M. R. Setare and F. Darabi, Gen. Rel. Grav. **44**, 2521 (2012).

[35] M. Cataldo, P. Mella, P. Minning and J. Saavedra, Phys. Lett. B **662**, 314 (2008).

[36] B. Gumjudpai and K. Thepsuriya, to appear in Astrophys. Space Sci. (2012), arXiv:1207.2920 [astro-ph.CO]

[37] B. Gumjudpai and C. Kaeonikhom, arXiv:1201.3499 [astro-ph.CO].

[38] C. Kaeonikhom, B. Gumjudpai and E. N. Saridakis, Phys. Lett. B **695**, 45 (2011).

[39] Y.-H. Wei, astro-ph/0405368.

[40] M. Kaplinghat, G. Steigman, I. Tkachev and T. P. Walker, Phys. Rev. D **59**, 043514 (1999).

[41] M. Kaplinghat, G. Steigman and T. P. Walker, Phys. Rev. D **61**, 103507 (2000).

[42] R. R. Caldwell, Phys. Lett. B **545**, 23 (2002); R. R. Caldwell, M. Kamionkowski and N. N. Weinberg, Phys. Rev. Lett. **91**, 071301 (2003).

[43] A. Dev, D. Jain and D. Lohiya, arXiv:0804.3491 [astro-ph].

[44] G. Sethi, A. Dev and D. Jain, Phys. Lett. B **624**, 135 (2005).

[45] S. Kumar, Mon. Not. Roy. Astron. Soc. **422**, 2532 (2012).

[46] D. Jain, A. Dev and J. S. Alcaniz, Class. Quan. Grav. **20**, 4163 (2003).

[47] Z. H. Zhu, M. Hu, J. S. Alcaniz and Y. X. Liu, Astron. and Astrophys. **483**, 15 (2008).

[48] S. W. Allen, R. W. Schmidt, A. C. Fabian, Mon. Not. Roy. Astro. Soc. **334**, L11 (2002); S. W. Allen, R. W. Schmidt, A. C. Fabian, H. Ebeling, Mon. Not. Roy. Astro. Soc. **342**, 287 (2003).